

Datasheet: MCA1846T

Description:	HAMSTER ANTI MOUSE CD81
Specificity:	CD81
Other names:	TAPA-1
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	Eat2
Isotype:	IgG1
Quantity:	25 µg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/100
Immunohistology - Frozen (1)	▪			
Immunohistology - Paraffin		▪		
ELISA	▪			
Immunoprecipitation	▪			
Western Blotting (2)	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

(1)The epitope recognised by this antibody is reported to be sensitive to formaldehyde fixation and tissue processing. Bio-Rad recommends the use of acetone fixation for frozen sections.

(2)Clone Eat2 recognizes mouse CD81 under non-reducing conditions.

Target Species	Mouse
Species Cross Reactivity	Reacts with: Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide

Carrier Free	Yes
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	38C13, murine B cell line.
External Database Links	<p>UniProt: P35762 Related reagents</p> <p>Entrez Gene: 12520 Cd81 Related reagents</p>
Synonyms	Tapa1
Fusion Partners	Spleen cells from immunised Armenian hamsters were fused with cells of the mouse PX3-Ag.8.653 myeloma cell line.
Specificity	<p>Hamster anti Mouse CD81 antibody, clone Eat2 recognizes mouse and rat CD81, also known as TAPA-1 or Target of the antiproliferative antibody 1. CD81 is a 236 amino acid ~26 kDa multipass transmembrane protein belonging to the TM4SF family (UniProt: P35762). In rodents CD81 is expressed at much higher levels on resting B cells than on T cells, although increased expression on T cells is found following activation. Hamster anti Mouse CD81 antibody, clone Eat2 induces homotypic aggregation of B cells and inhibits anti Ig and IL-4 induced proliferation (Maecker et al. 2000). Eat 2 requires the presence of both extracellular loops of TAPA-1 for binding.</p> <p>Mice lacking CD81 demonstrate reduced fertility through impaired oocyte-sperm fusion, double knockout CD81^{-/-} CD9^{-/-} mice are completely infertile suggesting complimentary roles in oocyte-sperm fusion (Rubenstein et al. 2006).</p>
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> Clark, K.L. <i>et al.</i> (2001) PGRL is a major CD81-associated protein on lymphocytes and distinguishes a new family of cell surface proteins. J Immunol. 167 (9): 5115-21. Maecker, H.T. <i>et al.</i> (2000) Differential expression of murine CD81 highlighted by new anti-mouse CD81 monoclonal antibodies. Hybridoma 19: 15-22. Conde-Vancells, J. <i>et al.</i> (2010) Candidate biomarkers in exosome-like vesicles purified from rat and mouse urine samples. Proteomics Clin Appl. 4 (4): 416-25. Conde-Vancells, J. <i>et al.</i> (2008) Characterization and comprehensive proteome profiling of exosomes secreted by hepatocytes. J Proteome Res. 7: 5157-66. Takeda, Y. <i>et al.</i> (2008) Double deficiency of tetraspanins CD9 and CD81 alters cell motility and protease production of macrophages and causes chronic obstructive pulmonary disease-like phenotype in mice. J Biol Chem. 283: 26089-97. Suzuki, M. <i>et al.</i> (2009) Tetraspanin CD9 negatively regulates lipopolysaccharide-induced macrophage activation and lung inflammation. J Immunol. 182: 6485-93. Ha, C.T. <i>et al.</i> (2005) Binding of pregnancy-specific glycoprotein 17 to CD9 on macrophages induces secretion of IL-10, IL-6, PGE2, and TGF-beta1. J Leukoc Biol. 77: 948-57. Pan, Q. <i>et al.</i> (2011) Hepatic cell-to-cell transmission of small silencing RNA can extend the therapeutic reach of RNA interference (RNAi). Gut. 61: 1330-9. Jin, Y. <i>et al.</i> (2013) Statins decrease lung inflammation in mice by upregulating tetraspanin CD9 in macrophages. PLoS One. 8: e73706. Royo, F. <i>et al.</i> (2013) Transcriptome of extracellular vesicles released by hepatocytes. PLoS

[One. 8: e68693.](#)

11. Owens, D.M. and Watt, F.M. (2001) Influence of beta1 integrins on epidermal squamous cell carcinoma formation in a transgenic mouse model: alpha3beta1, but not alpha2beta1, suppresses malignant conversion. [Cancer Res. 61: 5248-54.](#)

12. Jin, Y. *et al.* (2018) Double deletion of tetraspanins CD9 and CD81 in mice leads to a syndrome resembling accelerated aging. [Sci Rep. 8 \(1\): 5145.](#)

Storage Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Shelf Life 18 months from date of despatch.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: 10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Hamster IgG (STAR104...) [DyLight®549](#), [DyLight®649](#), [DyLight®800](#),
[FITC](#)

Goat Anti Hamster IgG (STAR79...) [Biotin](#), [FITC](#), [HRP](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M315411:180503'

Printed on 20 Jun 2018

© 2018 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)