Datasheet: MCA342PE

Description:	MOUSE ANTI RAT CD163:RPE
Specificity:	CD163
Other names:	ED2
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	ED2
Isotype:	IgG1
Quantity:	100 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Rat

Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

Reconstitution

Reconstitute with 1 ml distilled water

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE 488nm laser	496	578

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernanant.

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide
1% Bovine Serum Albumin
5% Sucrose

Immunogen

Rat Spleen cell homogenate.

Fusion Partners

Spleen cells from immunised BALB/c mice were fused with cells of the SP2/0-Ag 14 mouse myeloma cell line.

Specificity

Mouse anti Rat CD163, clone ED2 recognises the rat ED2 cell surface glycoprotein (Dijkstra *et al.* 1985). A 175 kDa molecule also known as rat CD163, a member of the group B scavenger receptor cysteine-rich (SRCR) family and an erythroblast adhesion receptor (Fabriek *et al.* 2007).

Mouse anti rat CD163, clone ED2 was shown to detect approximately 50% of peritoneal macrophages, a subset of splenic macrophages, and most tissue macrophages. However, no staining was observed in monocytes or alveolar macrophages (Dijkstra *et al.* 1985, Beelen *et al.* 1987). In freshly isolated bone marrow, expression of CD163 was limited to mature macrophages only (Barbe *et al.* 1990).

Clone ED2 may be used in immunohistology using antigen retrieval, and has also been described reacting with paraffin-embedded material following PLP fixation (Periodate-lysine-paraformaldehyde), see Whiteland *et al.*

Flow Cytometry

Use 10ul of the suggested working dilution to label 10^6 cells in 100ul.

References

1. Dijkstra, C.D. *et al.* (1985) The heterogeneity of mononuclear phagocytes in lymphoid organs: distinct macrophage subpopulations in the rat recognized by monoclonal antibodies ED1, ED2 and ED3. Immunology. 54 (3): 589-99.
2. Beelen, R.H.J. *et al.* (1987) Monoclonal antibodies ED1, ED2 and ED3 against rat macrophages: Expression of recognized antigens in different stages of differentiation. Transplant Proc. 3: 3166-70.
3. Barbe, E. *et al.* (1990) Characterization and expression of the antigen present on resident rat macrophages recognized by monoclonal antibody ED2. Immunobiol. 182: 88-99.
4. Dijkstra, C.D. & Damoiseaux, J.G. (1993) Macrophage heterogeneity established by immunocytochemistry. Prog Histochem Cytochem. 27 (2): 1-65.
5. Whiteland, J.L. *et al.* (1995) Immunohistochemical detection of T-cell subsets and other leukocytes in paraffin-embedded rat and mouse tissues with monoclonal antibodies. J Histochem Cytochem. 43 (3): 313-20.
6. Polfliet, M.M.J. *et al.* (2002) Identification of the rat mature macrophage antigen ED2 as CD163: Regulation by glucocorticoids and role in the production of proinflammatory mediators. PhD Thesis. Vrije University, Amsterdam.
7. Deng, X. *et al.* (2005) Chronic alcohol consumption accelerates fibrosis in response to cerulein-induced pancreatitis in rats. Am J Pathol. 166 (1): 93-106.
8. Piquet-Pellorce, I. *et al.* (2005) Identification of the leukemia inhibitory factor cell targets within the rat testis. Biol Reprod. 72: 602-11.
9. Fujita, E. *et al.* (2010) Statin attenuates experimental anti-glomerular basement membrane glomerulonephritis together with the augmentation of alternatively activated macrophages. Am J Pathol. 177 (3): 1143-54.
10. Schwartzkopff, J. *et al.* (2010) NK cell depletion delays corneal allograft rejection in baby rats. Mol Vis. 16: 1928-35.
11. Baker, S.C. *et al.* (2011) Cellular integration and vascularisation promoted by a resorbable, particulate-leached, cross-linked poly(ε-caprolactone) scaffold. Macromol Biosci. 11 (5): 618-27.
12. Bedi, A. *et al.* (2010) Effect of early and delayed mechanical loading on tendon-to-bone healing after anterior cruciate ligament reconstruction. J Bone Joint Surg Am. 92: 2387-401.
13. Banerjee, S. *et al.* (2003) Development of organised conjunctival leucocyte aggregates after corneal transplantation in rats. Br J Ophthalmol. 87: 1515-22.
14. Moghaddami, M. *et al.* (2005) MHC class II compartment, endocytosis and phagocytic activity of macrophages and putative dendritic cells isolated from normal tissues rich in synovium. Int Immunol. 17: 1117-30.
15. Wojcik, M. *et al.* (2012) Immunodetection of cyclooxygenase-2 (COX-2) is restricted to tissue macrophages in normal rat liver and to recruited mononuclear phagocytes in liver injury and cholangiocarcinoma. Histochem Cell Biol. 137: 217-33.
16. Kawakami, A.P. *et al.* (2011) Inflammatory Process Modulation by Homeopathic Arnica montana

6CH: The Role of Individual Variation. Evid Based Complement Alternat Med. 2011: 917541.
17. Kajita, M. *et al.* (2011) iNOS expression in vascular resident macrophages contributes to circulatory dysfunction of splanchnic vascular smooth muscle contractions in portal hypertensive rats. Am J Physiol Heart Circ Physiol. 300: H1021-31.
18. Jiao, K. *et al.* (2013) The Identification of CD163 Expressing Phagocytic Chondrocytes in Joint Cartilage and Its Novel Scavenger Role in Cartilage Degradation. PLoS One. 8(1):e53312.
19. Fujii, Y. *et al.* (2013) Effect of enzymatically modified isoquercitrin on preneoplastic liver cell lesions induced by thioacetamide promotion in a two-stage hepatocarcinogenesis model using rats. Toxicology. 305: 30-40.
20. Lobato-Pascual, A. *et al.* (2013) Rat macrophage C-type lectin is an activating receptor expressed by phagocytic cells. PLoS One. 8: e57406.
21. Fabriek, B.O. *et al.* (2007) The macrophage CD163 surface glycoprotein is an erythroblast adhesion receptor. Blood. 109 (12): 5223-9.
22. Park, E.S. *et al.* (2014) Establishment of a rat model for canine necrotizing meningoencephalitis (NME). Vet Pathol. 51 (6): 1151-64.
23. Keitel, V. *et al.* (2008) Expression and function of the bile acid receptor TGR5 in Kupffer cells. Biochem Biophys Res Commun. 372: 78-84.
24. Hozumi, Y. *et al.* (2015) Expression and localization of the diacylglycerol kinase family and of phosphoinositide signaling molecules in adrenal gland. Cell Tissue Res. 362 (2): 295-305.
25. Fernandez-Bustamante, A. *et al.* (2015) Brief Glutamine Pretreatment Increases Alveolar Macrophage CD163/Heme Oxygenase-1/p38-MAPK Dephosphorylation Pathway and Decreases Capillary Damage but Not Neutrophil Recruitment in IL-1/LPS-Insufflated Rats. PLoS One. 10 (7): e0130764.
26. Adamo, H.H. *et al.* (2015) Adaptive (TINT) Changes in the Tumor Bearing Organ Are Related to Prostate Tumor Size and Aggressiveness. PLoS One. 10 (11): e0141601.
27. Santos, L. *et al.* (2016) *In vitro* and *in vivo* assessment of magnetically actuated biomaterials and prospects in tendon healing. Nanomedicine (Lond). 11 (9): 1107-22.
28. Ibarra V *et al.* (2016) Evaluation of the Tissue Response to Alginate Encapsulated Islets in an Omentum Pouch Model. J Biomed Mater Res A. May 3. [Epub ahead of print]
29. Tentillier N *et al.* (2016) Anti-Inflammatory Modulation of Microglia via CD163-Targeted Glucocorticoids Protects Dopaminergic Neurons in the 6-OHDA Parkinson's Disease Model. J Neurosci. 36 (36): 9375-90.
30. Zakrzewicz, A. *et al.* (2015) Monocytic Tissue Transglutaminase in a Rat Model for Reversible Acute Rejection and Chronic Renal Allograft Injury. Mediators Inflamm. 2015: 429653.
31. Rave-Fränk M *et al.* (2013) Rat model of fractionated (2 Gy/day) 60 Gy irradiation of the liver: long-term effects. Radiat Environ Biophys. 52 (3): 321-38.
32. Wang, M. *et al.* (2017) Characterization of the Micro-Environment of the Testis that Shapes the Phenotype and Function of Testicular Macrophages. J Immunol. May 1. pii: 1700162. [Epub ahead of print]
33. Stavenuiter, A.W. *et al.* (2015) Protective Effects of Paricalcitol on Peritoneal Remodeling during Peritoneal Dialysis. Biomed Res Int. 2015: 468574.
34. Hawkins, K.E. *et al.* (2017) Targeting resolution of neuroinflammation after ischemic stroke with a lipoxin A_4 analog: Protective mechanisms and long-term effects on neurological recovery. Brain Behav. 7 (5): e00688.
35. Almahrog, A.J. *et al.* (2016) In vivo association of immunophenotyped macrophages expressing CD163 with PDGF-B in gingival overgrowth-induced by three different categories of medications. J Oral Biol Craniofac Res. 6 (1): 10-7.
36. Han, T.T. *et al.* (2015) Adipose-derived stromal cells mediate in vivo adipogenesis, angiogenesis and inflammation in decellularized adipose tissue bioscaffolds. Biomaterials. 72: 125-37.
37. Pannell, M. *et al.* (2016) Adoptive transfer of M2 macrophages reduces neuropathic pain via opioid peptides. J Neuroinflammation. 13 (1): 262.

Storage	Prior to reconstitution store at +4ºC. Following reconstitution store at +4ºC. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA1209PE)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

'M300672:170106'

Printed on 25 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint