Datasheet: MCA4633PET

Description:	MOUSE ANTI RAT CD106:RPE
Specificity:	CD106
Other names:	VCAM-1
Format:	RPE
Product Type:	Monoclonal Antibody
Clone:	MR106
Isotype:	IgG1
Quantity:	25 TESTS

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Rat

Product Form

Purified IgG conjugated to R. Phycoerythrin (RPE) - lyophilized

Reconstitution

Reconstitute with 0.25ml distilled water

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
RPE 488nm laser	496	578

Preparation

Purified IgG prepared by affinity chromatography on Protein G from tissue culture supernatant

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

0.09% Sodium Azide (NaN_3)
1% Bovine Serum Albumin
5% Sucrose

Immunogen

Rat CD106-transfected L5178Y cells.

External Database Links

UniProt:
P29534 Related reagents

	Entrez Gene: 25361 Vcam1 Related reagents
Synonyms	Vcam-1
Specificity	**Mouse anti Rat CD106 antibody, clone MR106** recognizes rat CD106, otherwise known as VCAM-1 (vascular adhesion molecule 1), a 110kDa inducible type I transmembrane glycoprotein and member of the immunoglobulin supergene family, which is predominantly expressed on vascular endothelium, and also on bone marrow stromal cells, follicular dendritic cells and some macrophages. CD106 interacts with the lymphocyte homing receptor VLA-4 (alpha4 beta1 integrin) and LPAM-1 (alpha4 beta7 integrin) and mediates leucocyte-endothelial cell adhesion and signal transduction. Expression of CD106 is upregulated following cytokine activation, and endothelial CD106 plays a role in the extravasation of leucocytes from blood vessels during inflammation.
Flow Cytometry	Use 10ul of the suggested working dilution to label $1x10^6$ cells in 100ul.
References	1. Miyao, N. *et al.* (2006) Various adhesion molecules impair microvascular leukocyte kinetics in ventilator-induced lung injury. Am J Physiol Lung Cell Mol Physiol. 290 (6): L1059-68. 2. Kubota, H. *et al.* (2007) Identification and characterization of vitamin A-storing cells in fetal liver: implications for functional importance of hepatic stellate cells in liver development and hematopoiesis. Stem Cells. 25 (9): 2339-49.
Further Reading	1. Bevilacqua, M.P. (1993) Endothelial-leukocyte adhesion molecules. Annu Rev Immunol. 11: 767-804.
Storage	Prior to reconstitution store at +4°C. After reconstitution store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.
Shelf Life	12 months from date of reconstitution.
Health And Safety Information	Material Safety Datasheet documentation #10075 available at: 10075: https://www.bio-rad-antibodies.com/uploads/MSDS/10075.pdf
Regulatory	For research purposes only

Related Products

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL:RPE (MCA1209PE)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

'M300596:170106'

Printed on 14 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint