Datasheet: VPA00437KT

Description:	ARS2 ANTIBODY WITH CONTROL LYSATE
Specificity:	ARS2
Format:	Purified
Product Type:	PrecisionAb™ Polyclonal
Isotype:	Polyclonal IgG
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively validated for the western blot application. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	20µl Goat polyclonal antibody purified by affinity chromatography
Buffer Solution	TRIS buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN_3) 0.5% Bovine Serum Albumin
Immunogen	Peptide sequence C-DPRAIVEYRDLDAPD
External Database Links	**UniProt:** Q9BXP5 Related reagents **Entrez Gene:** 51593 SRRT Related reagents
Synonyms	ARS2, ASR2
Specificity	**Goat anti Human ARS2 antibody** recognizes ARS2 also known as arsenate resistance protein 2, serrate RNA effector molecule homolog or SRRT.

	Goat anti Human ARS2 antibody detects a band of 117 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti ARS2 detects a band of approximately 117 kDa in Jurkat cell lysates
Instructions For Use	Please refer to the PrecisionAb western blotting protocol. For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400µg Jurkat lysate lyophilized in RIPA buffer.
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190µl DI H_2O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. - If using BME reconstitute the lyophilized lysate with 180µl DI H_2O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20ºC, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Shelf Life	As supplied, 12 months from date of despatch
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories
Health And Safety Information	Material Safety Datasheet documentation #10058 #10561 available at: Antibody (10058): https://www.bio-rad-antibodies.com/uploads/MSDS/10058.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Donkey Anti Sheep IgG (STAR88...) HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	**Worldwide**	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	**Europe**	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

'M295393:161101'

Printed on 01 May 2018

© 2018 Bio-Rad Laboratories Inc | Legal | Imprint